

Endocrine Disruptors

ISSN: (Print) 2327-3747 (Online) Journal homepage: <http://tandfonline.com/loi/kend20>

Comparison of vehicle mortality following in ovo exposure of Japanese quail (*Coturnix japonica*) eggs to corn oil, triolein and a fatty acid mix

Karen M. Dean, Leah D. Baltos, Tiffany Carro, Andrew, N. Iwaniuk, Meredith

E.B. Bohannon & Mary Ann Ottinger

To cite this article: Karen M. Dean, Leah D. Baltos, Tiffany Carro, Andrew, N. Iwaniuk, Meredith

E.B. Bohannon & Mary Ann Ottinger (2016): Comparison of vehicle mortality following in ovo exposure of Japanese quail (*Coturnix japonica*) eggs to corn oil, triolein and a fatty acid mix,

Endocrine Disruptors, DOI: 10.1080/23273747.2016.1224022

To link to this article: <http://dx.doi.org/10.1080/23273747.2016.1224022>

© 2016 Taylor & Francis Group, LLC

Accepted author version posted online: 22

Aug 2016.

Published online: 22 Aug 2016.

Submit your article to this journal

Article views: 12

View related articles

View Crossmark data

Comparison of vehicle mortality following *in ovo* exposure of Japanese quail (*Coturnix japonica*) eggs to corn oil, triolein and a fatty acid mix

Karen M. Dean^{1,2}, Leah D. Baltos², Tiffany Carro², Andrew, N. Iwaniuk³, Meredith E.B.

Bohannon² and Mary Ann Ottinger^{2,4,*}

¹ Abt Associates, Boulder CO

2 Department of Animal and Avian Sciences, University of Maryland, College Park;

3 Department of Neuroscience, University of Lethbridge, Lethbridge, Alberta, Canada

4 Department of Biology and Biochemistry, University of Houston, Houston, Texas 77204

*Corresponding author: Mary Ann Ottinger (maotting@central.uh.edu)

ABSTRACT

The use of avian egg bioassays for the determination of embryonic mortality and development effects of toxicant exposure is widespread in ecotoxicology. While these studies have a number of experimental limitations to consider, they offer a rapid, cost effective alternative to maternal feeding studies. In preparing to conduct such studies a number of factors must be taken into consideration, including solubility of the toxicant, dissolution solvent, injection site, volume and incubation position. Species-specific requirements for humidity and position should be considered in order to optimize successful incubation with different species. Japanese quail eggs, were injected prior to incubation with 1 μ l or 5 μ l of corn oil, triolein or a fatty acid mix, using air cell or albumen injection. Eggs were incubated according to standard poultry practices or in a prone position to determine if there were any differences in hatching success. Hatching success was reduced in eggs that were injected with 5 μ l and those incubated in a prone position. The highest rate of hatching success was observed for eggs injected with 1 μ l of the fatty acid mixture through the air cell.

Keywords

embryonic mortality, Japanese quail, vehicle mortality, incubation position, air cell injections, albumen injections